

Preprint IFUNAM FT 94-43

March 1994

May 1994 (corrected)

Two electrons in an external oscillator potential: hidden algebraic structure

by

Alexander Turbiner[†]

Instituto de Fisica, UNAM, Apartado Postal 20-364, 01000 Mexico D.F.,
Mexico

ABSTRACT

It is shown that the Coulomb correlation problem for a system of two electrons (two charged particles) in an external oscillator potential possesses a hidden sl_2 -algebraic structure being one of recently-discovered quasi-exactly-solvable problems. The origin of existing exact solutions to this problem, recently discovered by several authors, is explained. A degeneracy of energies in electron-electron and electron-positron correlation problems is found. It manifests the first appearance of hidden sl_2 -algebraic structure in atomic physics.

PASC: 03.65.Fd and 03.65.Ge

[†]On leave of absence from the Institute for Theoretical and Experimental Physics, Moscow 117259, Russia

E-mail: turbiner@teorica0.ifisicacu.unam.mx or turbiner@vxcern.cern.ch

The problem of evaluation of effects of inter-electronic interactions is one of the central problems in atomic physics. The main difficulty comes out from the fact that this problem cannot be solved exactly even in particular cases, while numerical solutions are too complicated to gain a proper intuition. Therefore, it is quite important to find and elaborate situations, where this problem can be modelled in some relevant way, admitting exact, analytic solutions. One of such situations has been described recently in [1, 2]. A system of two electrons in an external harmonic-oscillator potential with an additional linear interaction was studied in the relative coordinate, defined by the Hamiltonian ¹

$$H = -\nabla_1^2 + \omega^2 r_1^2 - \nabla_2^2 + \omega^2 r_2^2 + \frac{2\beta}{|\mathbf{r}_1 - \mathbf{r}_2|} + \lambda|\mathbf{r}_1 - \mathbf{r}_2| \quad (1)$$

where $r_{1,2}$ are the coordinates of the electrons and $\beta = 1$. Atomic units $\hbar = m = e = 1$ are used throughout and an overall factor $\frac{1}{2}$ is omitted. It was found that for certain values of oscillator frequency ω and the parameter λ some eigenstate of (1) can be obtained analytically. The main purpose of this note is to show that this feature is nothing but a consequence of the fact that (1) is one of recently-discovered quasi-exactly-solvable Schroedinger operators [4, 5]. It implies an existence of a hidden algebraic structure [6]. Hereafter, we will focus on the case $\lambda = 0$.

Quasi-exactly-solvable problems are quantum-mechanical problems for which several eigenstates can be found explicitly. They occupy an intermediate place between exactly-solvable (like Coulomb potential, harmonic oscillator etc) and non-solvable. The quasi-exactly-solvable Schroedinger equations appear in two forms: (i) the Hamiltonian with an infinite discrete spectrum with several eigenstates known algebraically and (ii) the Hamiltonian depending on a free parameter, say β , and a certain fixed magnitude of energy corresponds to the i th state of the Hamiltonian at i th value of parameter β (where $i = 0, 1, 2, \dots, n$) ². Those problems are named the first- and the second type, respectively. Surprisingly, exactly-solvable problems

¹ In [3] so called pseudo atoms were introduced : quantum system made from atomic ones in which all Coulomb interactions are replaced by oscillator ones, attractive or repulsive as the case may be. The system described by (1) can be treated as modified two-electron pseudo atom: where Coulomb attractive interactions are replaced by oscillator ones, while Coulomb repulsion remains unchanged or modified by the linear interaction.

²Precisely speaking, this means that for the parameters $\beta_0, \beta_1, \dots, \beta_n$, the ground-

like the Coulomb problem, the Morse oscillator, the Pöschl-Teller potential have two equivalent representations, either as the first-type problems or as the second-type ones [4]³.

The underlying idea behind quasi-exactly-solvability is the existence of a hidden algebraic structure. Let us recall a general construction considering the one-dimensional Schroedinger equation as an example. Take the algebra sl_2 realized in the first-order differential operators

$$\begin{aligned} J_n^+ &= r^2 d_r - nr, \\ J_n^0 &= r d_r - \frac{n}{2}, \\ J_n^- &= d_r, \end{aligned} \quad (2)$$

where $r \in R$ and $d_r \equiv \frac{d}{dr}$. Those three generators obey sl_2 -algebra commutation relations for any value of the parameter n . If n is a non-negative integer, the algebra (2) possesses $(n+1)$ -dimensional irreducible representation

$$\mathcal{P}_{n+1}(r) = \langle 1, r, r^2, \dots, r^n \rangle. \quad (3)$$

It is evident, that taking any polynomial in the generators (2), we arrive at a differential operator having the space (3) as the finite-dimensional invariant subspace. In other words, almost any polynomial in the generators (2) possesses $(n+1)$ eigenfunctions in the form of polynomial in r of degree n .

Let us take the quasi-exactly-solvable operator⁴

$$T_2 = -J_n^0 J_n^- + 2\omega_r J_n^+ - (n/2 + 2l + 2)J_n^- - \omega_r n. \quad (4)$$

Substituting (2) into (4), one gets the differential operator

$$T_2(r, d_r; n) = -r d_r^2 + 2(\omega_r r^2 - l - 1)d_r - 2\omega_r n r \quad (5)$$

for which one can define the spectral problem

$$T_2(r, d_r; n)p(r) = -\beta(n)p(r) \quad (6)$$

state energy at β_0 is equal to the energy of the first-excited state at β_1 , is equal to the energy of 2th-excited state at β_2 etc, is equal to the energy of n th excited state at β_n .

³For the Coulomb the second-type representation is nothing but the well-known Sturm representation.

⁴It belongs to the case VIII in the classification [5].

where $\beta(n)$ is a spectral parameter. It is clear, that this problem possesses $(n+1)$ eigenfunctions, $p_0(r), p_1(r), p_2(r), \dots, p_n(r)$ in the form of polynomial of the n -th power. Other eigenfunctions are non-polynomial and in general, they cannot be found in closed analytic form. Now let us make a gauge transformation in (5),(6), introducing a new function,

$$u(r) = r^{l+1} p(r) \exp(-\sqrt{\omega_r} r^2 / 2), \quad (7)$$

then make a replacement in the last term in (5):

$$2\omega_r n = \epsilon' - \omega_r(2l+3) \quad (8)$$

where ϵ' is a new parameter, and divide (6) over r . Finally, we obtain the equation

$$[-d_r^2 + \omega_r^2 r^2 + \frac{\beta}{r} + \frac{l(l+1)}{r^2}]u(r) = \epsilon' u(r) \quad (9)$$

Putting in (9) $\beta = 1$ and saying that now a new spectral parameter is ϵ' , we arrive at the equation (9) of the paper [2]. If $\omega_r = 2\omega$ and $2\epsilon' = \epsilon$ is the energy of the relative motion, this equation appears in [2] as a radial equation for the relative motion in (1) after separation of the c.m. motion.

Equation (9) is a particular case of the quasi-exactly-solvable Schroedinger equation of the second type (Case VIII in the classification [5])⁵. From the physical viewpoint, the parameter β in (1) has a meaning of the constant of the inter-electronic interaction. This parameter can be changed by replacing an electron by a charged particle with charge Z . In principle, keeping the frequency ω and the energy ϵ' fixed, for any n and l one can find $(n+1)$ systems of two particles with different charges in the oscillator potential related each other via hidden sl_2 algebraic structure (see a discussion in footnote 2).

Now let us describe some features of the eigenvalue problem (6).

- (i). It is clear that the operator T_2 is self-adjoint and hence its eigenvalues are real. The first $(n+1)$ eigenvalues coincide with the eigenvalues of the Jacobian matrix with vanishing diagonal matrix elements

⁵ General case VIII corresponds to $\lambda \neq 0$. In [4] this problem was named the *generalized Coulomb problem*.

$$\hat{H} = \begin{pmatrix} 0 & 2\omega_r & 0 & \cdots & 0 & 0 & \cdots \\ n(n+1+2l) & 0 & 4\omega_r & \cdots & 0 & 0 & \cdots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & \cdots & (i+1)(i+2+2l) & 0 & 2(n-i+1)\omega_r & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 & \cdots & 0 & 2n\omega_r \\ 0 & 0 & \cdots & 0 & \cdots & 2(l+1) & 0 \end{pmatrix} \quad (10)$$

One can show that the spectrum of (10) is symmetric :

$$S_{n+1}(\beta) = (-1)^n S_{n+1}(-\beta) , \quad S_{n+1}(\beta) \equiv \det ||\hat{H} - \beta|| \quad (11)$$

that follows from the fact that all odd powers of the matrix $\hat{H}$ are traceless, $tr \hat{H}^{2j+1} = 0, j = 0, 1, \dots$ ⁶. So, for any $\omega_r > 0$ there exist $[(n+1)/2]$ ⁷ positive eigenvalues and the same amount negative eigenvalues. This property leads to an important conclusion: for the fixed n, l and ω_r there exist two eigenstates, one at $\beta > 0$ and another at $\beta < 0$, degenerate in energy (see (8)). In particular, this may allow the electron-electron correlation energy to be related with the electron-positron one in the problem (1) (see discussion below).

- (ii). Another important property of (6) is that all eigenvalues $\beta^{(n)} \propto \sqrt{\omega_r}$ and, hence, depend on ω_r monotonously. For instance,

$$\beta_{\pm}^{(1)} = \pm 2\sqrt{\omega_r(l+1)}$$

$$\beta_{\pm,0}^{(2)} = \{\pm 2\sqrt{\omega_r(4l+5)}, 0\}$$

In order to find the eigenvalues of (6) it is enough to perform calculations in one point on ω_r -axe, e.g. at $\omega_r = 1$.

⁶I am grateful to P. Mello for a discussion of this point.

⁷ $[a]$ means integer part of a .

The situation becomes slightly more complicated, if we want to keep the parameter β fixed in the formula (1), declaring that now we want to consider namely two-electron (or electron-positron) system, which implies $\beta = 1(-1)$. The relevant formulation of the problem is the following.

Let us fix n and l . This defines unambiguously the functional form of the pre-exponential factor in (7). Take a positive eigenvalue β in (6), which corresponds to repulsion of the particles in (1)). It depends on the parameter ω_r monotonously, growing from zero up to infinity, which means that one can always find the value of ω_r for what $\beta = 1$ or any positive number. Since there exist $[(n+1)/2]$ positive eigenvalues of β (see above), each of them is equal to one for a certain value of ω_r . Correspondingly, the lowest eigenvalue (ground state, no nodes in $p(r)$ following the oscillation theorem) leads to the smallest value of ω_r , the next eigenvalue leads to bigger value of ω_r (one positive root in $p(r)$) etc. Finally, we arrive at $[(n+1)/2]$ values of the parameter ω_r , for each of them the problem (1) has the analytic solution of the form (7) with $p(r)$ as a polynomial of the n th degree with number of positive roots varying from 0 up to $[(n+1)/2]$ (see Fig.1, where the case $n = 3$ is described as an illustrative example)⁸.

Taking the negative eigenvalues of β in (1) (that corresponds to attraction of the particles), one can repeat above considerations with the only difference that the number of positive roots varies from $[(n+1)/2]$ up to n . Following the property (i) for any eigenstate from algebraized part of the spectra (see above) of the problem (1) with positive β one can find an eigenstate with negative β with the same energy. For example, for the fixed n and the minimal ω_r : $\omega_r^{(0)}$, the ground state energy at $\beta = 1$ ($p(r)$ has no positive roots) is equal to the the energy of the n th excited state at $\beta = -1$ ($p(r)$ has n positive roots). For $\omega_r^{(1)}$, the energy of the 1st excited state at $\beta = 1$ ($p(r)$ has one positive root) is equal to the energy of the $(n-1)$ th excited state at $\beta = -1$ ($p(r)$ has $(n-1)$ positive roots) etc (see e.g. Fig.1). This is reminiscent of the situation in one-dimensional supersymmetric quantum mechanics by Nicolai-Witten, where if the supersymmetry is unbroken, all states of bosonic sector (except lowest one) are degenerate with the states of fermionic one.

⁸it explains a systematics found in numerical calculation in [2].

It is worth noting that recently it was shown [6] that whenever some analytic solutions for eigenfunctions of a certain one-dimensional (or reduced to one-dimensional) Schroedinger equation occur, it signals the existence of the hidden algebra sl_2 . Our present results manifest the first appearance of quasi-exactly-solvability in atomic physics. Novel developments of hidden algebra method in quantum mechanics, solid-state physics and quantum field theory, and also mathematical foundations can be found in Refs. [7].

I would like to thank C. Bunge for bringing my attention to the papers [1, 2] and useful comments. I am grateful to the Instituto de Fisica, UNAM for kind hospitality extended to me.

References

- [1] A. Samanta, S. Ghosh, *Phys. Rev.*, **A 42** , 1178 (1990)
- [2] M. Taut, *Phys. Rev.*, **A 48** , 3561 (1993)
- [3] M. Moshinsky, O. Novaro, A. Calles, *Journ. de Phys.*, **31** , C 4-125 (1970)
- [4] A.V. Turbiner, *Sov.Physics-ZhETF*, **94**, 33-44 (1988)
- [5] A.V. Turbiner, *Comm. Math. Phys.*, **118** , 467 (1988)
- [6] A.V. Turbiner, *Journ. Phys.*, **A25** , L1087 (1992)
- [7] A.Yu. Morozov, A.M. Perelomov, A.A. Rosly, M.A. Shifman and A.V. Turbiner, *Int.Journ.Mod. Phys.* **A5** , 803-843 (1990) ;
V.V. Ulyanov, O.B. Zaslavsky, *Phys. Reps* **216** ,179 (1992) ;
P.B. Wiegmann and A.V. Zabrodin, *J. Frank Institute Preprints*, Oct., Dec. 1993 (cond-mat/9310017, 9312088);
A.V. Turbiner, in “Lie Algebras, Cohomologies and New Findings in Quantum Mechanics”, *Contemporary Mathematics*, AMS, pp. 263-310 (1993), N. Kamran and P. Olver (Eds.)

This figure "fig1-1.png" is available in "png" format from:

<http://arXiv.org/ps/hep-th/9406018v1>